


Product no **AS21 4528****Anti-PEC1 | Plastid Envelope Channel 1****Product information**

Immunogen	The soluble domain 466 (M244 until stop codon, 64 kDa) was cloned into pET16b and transformed into BLR 21 for 467 expression in <i>Escherichia coli</i> UniProt: Q8VZM7-1 , TAIR: At5g02940
Host	Rabbit
Clonality	Polyclonal
Purity	Serum
Format	Lyophilized
Quantity	50 µl
Reconstitution	For reconstitution add 50 µl, of sterile water
Storage	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

Additional information | This antibody is recognizing PEC1, but not PEC2 or DMI1**Application information**

Recommended dilution	1 : 1000 (WB)
Expected apparent MW	100 75 kDa
Confirmed reactivity	<i>Arabidopsis thaliana</i>
Predicted reactivity	<i>Camelina sativa</i> , <i>Capsella rubella</i> , <i>Nicotiana benthamiana</i> , <i>Noccaea caerulea</i> , <i>Pisum sativum</i> , <i>Raphanus sativus</i> , Species of your interest not listed? Contact us
Selected references	Völkner et al (2021) Two plastid POLLUX ion channel-like proteins are required for stress-triggered stromal Ca ²⁺ -release, <i>Plant Physiology</i> , Volume 187, Issue 4, December 2021, Pages 2110–2125, https://doi.org/10.1093/plphys/kiab424



Samples:

- 1 – 2,5 µg of *Arabidopsis thaliana* mutant pec1-1pec2-1
- 2 – 2,5 µg of *Arabidopsis thaliana* mutant pec1-2pec2-2
- 3 – 2,5 µg of *Arabidopsis thaliana* whole leaf extract

Arabidopsis thaliana tissue was frozen with liquid nitrogen and ground to a fine powder with mortar and pestle. Protein was extracted by mixing with extraction buffer (200 mM tris pH 8.0, 4% (w/v) sodium dodecyl sulfate) to 0.5 g fresh weight/mL and heating at 80°C for 8 min. 5 µl protein extract were separated on 10 % SDS-PAGE and blotted at 75v for 45min to nitrocellulose (pore size of 20 µm), using wet transfer. Blot was blocked with 5% milk for 1h/RT with agitation. Blot was incubated in the primary antibody at a dilution of 1: 1000 ON/4°C in 5% Milk-TBS-T with agitation. The antibody solution was decanted and the blot was rinsed briefly twice, then washed once for 15 min and 3 times for 5 min in TBS-T at RT with agitation. Blot was incubated in matching secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated) diluted to 1:10 000 in for 1h/RT with agitation. The blot was washed as above and developed for 1 min with chemiluminescent detection reagent following manufacture's recommendations. Exposure time was 120 seconds.

Courtesy of Carsten Voelkner, Kunz Lab