


Product no **AS15 2863****Anti-Phly | DNA photolyase (At4g25290) (N-terminal part)****Product information**

Immunogen	KLH-conjugated peptide derived from <i>Arabidopsis thaliana</i> DNA photolyase, UniProt: F4JSJ6 , TAIR: AT4G25290 , located in the N-terminal part of the protein
Host	Rabbit
Clonality	Polyclonal
Purity	Immunogen affinity purified serum in PBS pH 7.4.
Format	Lyophilized
Quantity	50 µg
Reconstitution	For reconstitution add 50 µl of sterile water
Storage	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

Application information

Recommended dilution	1 : 1000 (WB)
Expected apparent MW	78 90 kDa
Confirmed reactivity	<i>Arabidopsis thaliana</i>
Not reactive in	No confirmed exceptions from predicted reactivity are currently known

application example

2,5 µg of total protein from *Arabidopsis thaliana* wild type darkness (1), wild type light (2) and insertion mutants: SALK_056328C darkness (3), SALK_056328C light (4), extracted with 0.1 M Tris-HCl pH 8.5, 4% SDS, 2% (v/v) 2-mercaptoethanol, 2 mM phenylmethylsulfonyl fluoride and denatured with Laemmli buffer at 95°C for 10 min were separated on 12% SDS-PAGE and blotted 2h to PVDF using semi-dry transfer. Blots were blocked with 5% milk PBS-T (Tween 0.5%) for 30 min. at room temperature (RT) with agitation. Blot was incubated in the primary antibody at a dilution of 1:1000 overnight at 4°C with agitation. The antibody solution was decanted and the blot was rinsed briefly twice, then washed 3 times for 10 min in 5% milk PBS-T at RT with agitation. Blot was incubated in secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated, from Agrisera) diluted to 1:25 000 in 5% milk PBS-T for 1h at RT with agitation. The blot was rinsed briefly twice, then washed 3 times for 10 min in PBS-T at RT with agitation. Blot was developed for 5 min with chemiluminescent detection reagent. Exposure time was 5 minutes.

Courtesy of Dr. Justyna Łabuz, Department of Biochemistry, Biophysics and Biotechnology, Jagiellonian University, Poland