

Product no **AS05 084****Anti-PsbA | D1 protein of PSII, C-terminal (rabbit antibody) (thylakoid membrane marker)****Product information**

Immunogen	KLH-conjugated synthetic peptide derived from available plant, algal and cyanobacterial PsbA sequences, including <i>Arabidopsis thaliana</i> UniProt: A4QJR4 , TAIR: AtCg00020 , <i>Oryza sativa</i> P0C434 , <i>Populus alba</i> Q14FH6 , <i>Physcomitrella patens</i> Q6YXN7 , <i>Chlamydomonas reinhardtii</i> P07753 , <i>Synechocystis</i> sp. P14660 and many others
Host	Rabbit
Clonality	Polyclonal
Purity	Serum
Format	Lyophilized
Quantity	50 µl
Reconstitution	For reconstitution add 50 µl of sterile water
Storage	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.
Additional information	<p>Due to biology of PsbA (D1) protein a number of degradation products can appear in a sample and may be observed when using anti-PsbA antibodies, including products having apparent molecular weights of 24 kDa and 16 kDa. D1 degradation is a complex set of events and the products observed can be influenced by both the extraction procedure and the physiology of the cells prior to harvest. Third, cross-linking may occur between D1 and cytochrome b559, shifting the protein higher in the gel. In cyanobacteria (PCC7942), three different bands were competed out by preincubating the antibody with the PsbA free peptide, indicating that all bands are indeed PsbA and its precursors or breakdown products. Competition assays were also performed with spinach and <i>Chlamydomonas</i>, confirming the identity of PsbA bands.</p> <p>Anti-PsbA antibodies will not detect D2 protein, as the peptide used to generate PsbA antibodies has no homology to the D2 sequence.</p> <p>This product can be sold containing ProClin if requested.</p>

Application information

Recommended dilution	1: 500 (IF), 1: 200 (IG), 1: 10 000 (WB)
Expected apparent MW	38 28-30 kDa
Confirmed reactivity	<p><i>Alniaria alnifolia</i>, <i>Anabaena</i> 7120, <i>Arabidopsis thaliana</i>, <i>Artemisia annua</i>, <i>Arundo</i> sp., <i>Begonia</i> sp., <i>Cannabis sativa</i> L., <i>Cladocopium goreau</i>, <i>Chlamydomonas reinhardtii</i>, <i>Chlorella ohadii</i>, <i>Chromera velia</i>, <i>Chlorella vulgaris</i>, <i>Colobanthus quitensis</i> (Kunth) Bartl, <i>Coscinodiscus wailesii</i>, <i>Craterostigma</i> sp., <i>Cyanidioschyzon merolae</i>, <i>Cytisus cantabricus</i> (Wilk.) Rchb. F, <i>Desmodium</i> sp., <i>Dianthus caryophyllus</i>, <i>Ditylum brightwellii</i>, <i>Eucalyptus globulus</i>, <i>Fraxinus rhynchophylla</i>, <i>Glycine max</i>, <i>Gongolaria barbata</i>, <i>Halomicronema hongdechloris</i>, <i>Hieracium pilosella</i> L., <i>Hordeum vulgare</i>, <i>Lasallia hispanica</i>, <i>Lindernia</i> sp., <i>Manihot esculenta</i>, <i>Marchantia polymorpha</i> (liverwort), <i>Medicago truncatula</i>, <i>Miscanthus x giganteus</i>, <i>Microcystis aeruginosa</i>, <i>Mirkania micrantha</i>, <i>Nicotiana benthamiana</i>, <i>Nicotiana tabacum</i>, <i>Panicum miliaceum</i>, <i>Panax ginseng</i>, <i>Panicum maximum</i>, <i>Paulinella chromatophora</i> (amoeba), <i>Pheodactylum tricornutum</i> CCAP 1055/1, <i>Physcomitrium patens</i>, <i>Picea glauca</i>, <i>Pinus strobus</i>, <i>Pisum sativum</i>, <i>Prochlorococcus</i> sp. (surface and deep water ecotype), <i>Salicornia bigelovii</i>, <i>Skeletonema costatum</i> (diatom), <i>Solanum lycopersicum</i>, <i>Spartina alterniflora</i>, <i>Spinacia oleracea</i>, <i>Spirodela polyrrhiza</i>, <i>Symbiodinium</i> sp., <i>Synechococcus</i> sp. PCC 7942, <i>Synechococcus elongatus</i> UTEX 2973, <i>Synechocystis</i> sp. 6803, <i>Syntrichia muralis</i>, <i>Thalassiosira weissflogii</i>, <i>Tetrademus obliquus</i>, <i>Triticum aestivum</i>, <i>Triticale</i>, <i>Zea mays</i>, <i>Quercus ilex</i></p>
Predicted reactivity	<p>Algae (brown and red), <i>Brassica napus</i>, <i>Carica papaya</i>, Conifers, Cyanobacteria, <i>Cannabis sativa</i>, <i>Citrus sinensis</i>, Dicots, <i>Eragrostis tef</i>, <i>Galdieria sulphuraria</i>, <i>Lactuca sativa</i>, <i>Lycopersicon esculentum</i>, <i>Medicago sativa</i>, <i>Nannochloropsis</i> sp., <i>Oryza sativa</i>, <i>Ostreococcus</i> sp., <i>Phalaenopsis aphrodite</i>, <i>Pisum sativum</i>, <i>Porphyridium purpureum</i>, <i>Sesamum indicum</i>, <i>Thalassiosira pseudonana</i>, <i>Zostera marina</i>, <i>Vitis vinifera</i> cellular [compartment marker] of thylakoid membrane</p> <p>Species of your interest not listed? Contact us</p>
Not reactive in	No confirmed exceptions from predicted reactivity are currently known
Additional information	The antibody is appropriate for detecting both, 24 kDa or the 10 kDa C-terminal fragments, whichever is generated under given treatment conditions. In our analysis we have seen both, ca. 24 kDa and ca. 10 kDa fragments from different samples, depending on treatments and isolation procedures.

Rabbit anti-PsbA antibody can detect more than one band of PsbA protein, e.g. precursor and mature protein as compare to the hen anti-PsbA antibodies AS01 016.


This antibody will detect the phosphorylated form of D1 as an alternate band to the main band on a high resolution gel.

The antibody will bind to cross-linked proteins: D1/D2, D1/cyt b559, D1/CP43.


The peptide is conserved in cyanobacterial D1:1 and D1:2.

Selected references

- [Popova](#) et al. (2025). Differential Photosynthetic Response of Tomato Plants—Ailsa Craig and Carotenoid Mutant tangerine—To Low Light Intensity and Low Temperature Treatment. *Crops* 2025, 5(6), 77; doi.org/10.3390/crops5060077.
- [Zhong](#) et al. (2025). Met246 and Asn250 in the D2 Protein are Essential for the Operation of the Quinone-Fe-Acceptor Complex of Photosystem II. *Plant Cell Physiol.* 2025 Jul 14;pcaf078. doi: 10.1093/pcp/pcaf078.
- [Lang](#) et al. (2025). Complexome profiling of the Chlamydomonas psb28 mutant reveals TEF5 as an early photosystem II assembly factor. *Plant Cell.* 2025 Jun 4;37(6):koaf055. doi: 10.1093/plcell/koaf055.
- [Konishi and Noguchi](#) (2025). The Mitochondrial Respiratory Chain is Important as an Electron Sink to Avoid Over-Reduction of Chloroplasts in Arabidopsis thaliana Leaves under High-Light Stress. *Plant Cell Physiol.* 2025 Jun 5;pcaf062. doi: 10.1093/pcp/pcaf062.
- [McKenzie and Puthiyaveetil](#) (2025). Protein phosphorylation and oxidative protein modification promote plant photosystem II disassembly for repair. *Plant Commun.* 2025 Mar 10;6(3):101202. doi: 10.1016/j.xplc.2024.101202.
- [Huang](#) et al. (2025). Nitrogen source type modulates heat stress response in coral symbiont (Cladocopium goreaui). *Appl Environ Microbiol.* 2025 Feb 19;91(2):e0059124. doi: 10.1128/aem.00591-24. Epub 2025 Jan 7.
- [Collombat](#) et al. (2025). Arabidopsis conditional photosynthesis mutants abc1k1 and var2 accumulate partially processed thylakoid preproteins and are defective in chloroplast biogenesis. *Commun Biol.* 2025 Jan 22;8(1):111. doi: 10.1038/s42003-025-07497-y.
- [Pica](#) et al. (2024). Functional ecological traits in young and adult thalli of canopy-forming brown macroalga *Gongolaria barbata* (Phaeophyta) from a transitional water system. *PeerJ.* 2024 Sep 12;12:e17959. doi: 10.7717/peerj.17959.
- [Salazar](#) et al. (2024). SOS1 tonoplast neo-localization and the RGG protein SALTY are important in the extreme salinity tolerance of *Salicornia bigelovii*. *Nat Commun.* 2024 May 20;15(1):4279. doi: 10.1038/s41467-024-48595-5.
- [Phukan](#) et al. (2024). Externally supplied ascorbic acid moderates detrimental effects of UV-C exposure in cyanobacteria. *Photochem Photobiol Sci.* 2024 Jul 12. doi: 10.1007/s43630-024-00612-8.
- [Zhao](#) et al. (2024). Psb28 protein is indispensable for stable accumulation of PSII core complexes in Arabidopsis. *Plant J.* 2024 May 26. doi: 10.1111/tj.16844.
- [Völkner](#) et al. (2024). Evidence for partial functional overlap of KEA and MSL transport proteins in the chloroplast inner envelope of *Arabidopsis thaliana*. *FEBS Lett.* 2024 Aug;598(15):1877-1887. doi: 10.1002/1873-3468.14887.



2 µg of total protein from (1) *Arabidopsis thaliana* leaf extracted with **Protein Extraction Buffer, PEB (AS08 300)**, (2) *Hordeum vulgare* leaf extracted with PEB, (3) *Chlamydomonas reinhardtii* total cell extracted with PEB, (4) *Synechococcus* sp. 7942 total cell extracted with PEB, (5) *Anabaena* sp. total cell extracted with PEB were separated on **4-12% NuPage** (Invitrogen) **LDS-PAGE** and blotted 1h to **PVDF**. Blots were blocked immediately following transfer in 2% ECL Advance blocking reagent (GE Healthcare) in 20 mM Tris, 137 mM sodium chloride pH 7.6 with 0.1% (v/v) Tween-20 (TBS-T) for 1h at room temperature with agitation. Blots were incubated in the primary antibody at a dilution of 1: 50 000 for 1h at room temperature with agitation. The antibody solution was decanted and the blot was rinsed briefly twice, then washed once for 15 min and 3 times for 5 min in TBS-T at room temperature with agitation. Blots were incubated in secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated, recommended secondary antibody **AS09 602**) diluted to 1:50 000 in 2% blocking solution for 1h at room temperature with agitation. The blots were washed as above and developed for 5 min with chemiluminescent detection reagent according the manufacturers instructions. Images of the blots were obtained using a CCD imager (FluorSMax, Bio-Rad) and Quantity One software (Bio-Rad).



Varying amounts of PsbA protein standard ([AS01 016S](#)) 250 fmol (**1**), 125 fmol (**2**), 62.5 fmol (**3**), 31.25 fmol (**4**), 15.625 fmol (**5**) and **2 µg of total protein** from Med4 (**6,7**) extracted with **Protein Extraction Buffer**, PEB ([AS08 300](#)). Samples were diluted with 1X sample buffer (NuPAGE LDS sample buffer (Invitrogen) supplemented with 50 mM DTT and heat at 70°C for 5 min and kept on ice before loading. Protein samples were separated on 4-12% Bolt Plus gels, LDS-PAGE and blotted for 70 minutes to PVDF using tank transfer. Blots were blocked immediately following transfer in 2% blocking reagent or 5% non-fat milk dissolved in 20 mM Tris, 137 mM sodium chloride pH 7.6 with 0.1% (v/v) Tween-20 (TBS-T) for 1h at room temperature with agitation. Blots were incubated in the primary antibody at a dilution of 1: 10 000 (in blocking reagent) for 1h at room temperature with agitation. The antibody solution was decanted and the blot was rinsed briefly twice, and then washed 1x15 min and 3x5 min with TBS-T at room temperature with agitation. Blots were incubated in secondary antibody (goat anti-rabbit IgG horse radish peroxidase conjugated, recommended secondary antibody [AS09 602](#), Agrisera) diluted to 1:25 000 in blocking reagent for 1h at room temperature with agitation. The blots were washed as above. The blot was developed with chemiluminescent detection reagent according the manufacturers instructions. Images of the blots were obtained using a CCD imager (VersaDoc MP 4000) and Quantity One software (Bio-Rad). Exposure time was 30 seconds.