

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

## Product no AS05 084A-HRP

## Anti-PsbA | D1 protein of PSII, C-terminal, HRP-conjugated (40 µg)

## **Product information**

Immunogen

KLH-conjugated synthetic peptide derived from available plant, algal and cyanobacterial PsbA sequences, including Arabidopsis thaliana UniProt: A4QJR4, TAIR: AtCg00020, Oryza sativa P0C434, Populus alba Q14FH6, Physcomitrella patens Q6YXN7, Chlamydomonas reinhardtii P07753, Synechocystis sp. P14660 and many others

**Host** Rabbit

Clonality Polyclonal

**Purity** Immunogen affinity purified serum in PBS pH 7.4, conjugated to HRP.

Format Liquid

Quantity 40 μg

Storage Store at 4°C for 12-18 months. A preservative may be added for long time storage up to 2 years.

Additional information

Due to biology of PsbA (D1) protein a number of degradation products can apprear in a sample and may be observed when using anti-PsbA antibodies, including products having apparent molecular weights of 24kDa and 16kDa. D1 degradation is a complex set of events and the products observed can be influenced by both the extraction procedure and the physiology of the cells prior to harvest. Third, cross-linking may occur between D1 and cytochrome b559, shifting the protein higher in the gel. In cyanobacteria (PCC7942), three different bands were competed out by preincubating the antibody with the PsbA free peptide, indicating that all bands are indeed PsbA and its precursors or breakdown products. Competition assays were also performed with spinach and Chlamydomonas, confirming the identity of PsbA bands.

Anti-PsbA antibodies will not detect D2 protein, as the peptide used to generate PsbA antibodies has no homology to the D2 sequence.

## **Application information**

Recommended dilution 1:15 000 (WB)

Expected | apparent

38 | 28-30 kDa

**Confirmed reactivity** 

Anabaena 7120, Arabidopsis thaliana, Artemisia annua, Arundo sp., Chlamydomonas reinhardtii, Colobanthus quitensis Kunt Bartl, Coscinodiscus wailesii, Craterostigma sp., Ditylum brightwellii, Glycine max, Hordeum vulgare, Lindernia sp., Miscanthus x giganteus, Marchantia polymorpha (liverwort), Nicotiana benthamiana, Panicum miliaceum, Panax ginseng, Panicum maximum, Pinus strobus, Physcomitrium patens, Synechococcus sp. PCC 7942, Paulinella chromatophora (amoeba), Prochlorococcus sp. (surface and deep water ecotype), Spirodela polyrhiza, Symbiodinium sp., Zea mays

Predicted reactivity

Algae (brown and red), Brassica napus, Conifers, Cyanobacteria, Dictos, Cannabis sativa, Galdieria sulphuraria, Lactuca sativa, Lycopersicum esculentum, Medicago sativa, Nannochloropsis sp., Oryza sativa, Ostreococcus sp. Pisum sativum, Porphyridium purpureum, Sesamum indicum, Thalassiosira pseudonana, Zosteria marina, Vitis vinifera cellular [compartment marker] of thylakoid membrane

Species of your interest not listed? Contact us

Not reactive in

No confirmed exceptions from predicted reactivity are currently known

Additional information

The antibody is appropriate for detecting both, 24 kDa or the 10 kDa C-terminal fragments, whichever is generated under given treatment conditions. In our analysis we have seen both, ca. 24 kDa and ca. 10 kDa fragments from different samples, depending on treatments and isolation procedures.

Rabbit anti-PsbA antibody can detect more than one band of PsbA protein, e.g. precursor and mature protein as compare to the hen anti-PsbA antibodies AS01 016.

This antibody will detect the phosphorylated form of D1 as an alternate band to the main band on a high resolution gel.

The antibody will bind to cross-linked proteins: D1/D2, D1/cyt b559, D1/CP43.

Selected references

Thurotte et al. (2020). DnaK3 Is Involved in Biogenesis and/or Maintenance of Thylakoid Membrane Protein Complexes in the Cyanobacterium Synechocystis Sp. PCC 6803. Life (Basel). 2020 Apr 30;10(5):E55. doi: 10.3390/life10050055.