

Product no **AS10 1003****Rabbit anti-Goat IgG Fc, HRP conjugated****Product information**

Host	Rabbit
Clonality	Polyclonal
Purity	Antigen Immunogen affinity purified rabbit IgG.
Format	Lyophilized
Quantity	1 mg
Reconstitution	For reconstitution add 1.1 ml of sterile water. Let it stand 30 minutes at room temperature to dissolve. Prepare fresh working dilutions daily
Storage	Store lyophilized material at 2-8°C. For long time storage after reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 1 mg of antibody in 1.1 ml of sterile water add 1.1 ml of glycerol. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well but without foaming.
Additional information	<p>This antibody reacts with the heavy chains on goat IgG based on immunoelectrophoresis.</p> <p>Minimum cross-reactivity is observed to the light chains or to non-immunoglobulin goat serum proteins based on immunoelectrophoresis.</p> <p>Antibody is supplied in 10 mM sodium phosphate, 150 mM sodium chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free and 0.1 % (v/v) Kathon CG is used as preservative. Use of sodium azide will inhibit enzymatic activity of horseradish peroxidase.</p> <p>BSA and milk have to be replaced by other blocking reagents, like donkey serum or commercial formulations which are free from bovine IgG.</p>

Application information**Recommended dilution** 1 : 500-1 : 5000 (IHC), 1 : 10 000-1 : 50 000 (WB)