

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no AS10 1281

Rabbit anti-Goat IgG Fc, ALP conjugated, min cross-reactivity to human serum Product information

Host Rabbit

Clonality Polyclonal

Purity Immunogen affinity purified IgG.

Format Liquid

Quantity 0.5 mg

Storage

Non-diluted antibody is stable for 4 years at 2-8 °C. For storage at -20 °C dilute antibody solution with an equal volume of glycerol to obtain final glycerol concentration of 50% to prevent loss of enzymatic activity. Such solution will not freeze in -20 °C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard, Be sure to mix well but without foaming.

Additional information

This antibody reacts with the heavy chains on goat IgG based on immunoelectrophoresis.

Minimum cross-reactivity is observed to the light chains or to non-immunoglobulin goat serum proteins based on immunoelectrophoresis. Minimum cross-reactivity is observed to human serum proteins based on immunoelectrophoresis.

Antibody is supplied in 30 mM triethanolamine, pH 7.2, 5 mM magnesium chloride, 0.1 mM zinc chloride, 1% (w/v) BSA, Protease/IgG free , 0.05% (w/v) sodium azide as preservative.

BSA and milk have to be replaced by other blocking reagents, like doneky serum or commercial formulations which are free from bovine IgG.

Application information

Recommended dilution 1:50-1:5000 (ICC), 1:20-1:2000 (IHC), 1:500-1:2000 (WB)