


Product no **AS19 4305****Anti-RPSA | 30S ribosomal protein S1 (bacterial)****Product information**

Immunogen	KLH-conjugated peptide derived from RPSA of <i>E.coli</i> , UniProt: P0AG67
Host	Rabbit
Clonality	Polyclonal
Purity	Immunogen affinity purified serum in PBS pH 7.4.
Format	Lyophilized
Quantity	50 µg
Reconstitution	For reconstitution add 50 µl, of sterile water
Storage	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

Application information

Recommended dilution	1: 100 (ELISA), 1: 3000 (WB)
Expected apparent MW	61 kDa
Confirmed reactivity	<i>Escherichia coli</i>
Predicted reactivity	bacteria, <i>Legionella pneumophila</i>
Not reactive in	No confirmed exceptions from predicted reactivity are currently known
Additional information	Antibody has been tested purified ribosomes from <i>E.coli</i> . Reactivity of this antibody in Western blot remains to be determined.
Selected references	To be added when available, antibody released in December 2020.



6 µg/well of total protein extracted freshly from *E. coli* strains MG1655, 536, UT189, RS218, IHE3034 and IHE3034ΔycgG2 with 20mM Tris buffer and extracted by freeze-thawing (3 cycles). The samples were then centrifuged for 30 min at 13 000 rpm and the protein content of the supernatants was subject to measuring with the BCA method. The proteins were separated on a 10 % SDS-PAGE and blotted onto a PVDF (pore size of 0,42 µm) membrane, using semi-dry transfer. The membrane was blocked with 5 % milk at 4°C/ON with agitation, followed by washing in PBS-T buffer (3 times for 15 min) before it was incubated with the primary antibody at a dilution of 1: 3 000 in PBS-T for 1h/RT with agitation. After that, the membrane was washed once for 30 min and 2 times for 15 min in PBST at RT with agitation. Then it was incubated in Agrisera matching secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated [AS09 602](#)) diluted 1:30 000 in PBS-T for 1h/RT with agitation. Finally, the immunoblot membrane was washed for 15 minutes three times with PBS-T and developed for 5 min with Agrisera ECLBright before imaging with a LAS 4000 image reader. Exposure time was 30 seconds.

Courtesy of Moa Näsman/the [Uhlen's Lab](#), Umeå University, Sweden