


Product no **AS21 4581****Anti-SnRK1α1 | SNF1-related protein kinase catalytic subunit alpha KIN10****Product information**

<b>Immunogen</b>	KLH-conjugated synthetic peptide derived from C-terminal part of <i>Arabidopsis thaliana</i> AKIN10 sequence UniProt: <a href="#">Q38997</a> , TAIR: <a href="#">At3g01090</a>
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Antigen affinity purified serum, in PBS pH 7.4
<b>Format</b>	Lyophilized
<b>Quantity</b>	50 µg
<b>Reconstitution</b>	For reconstitution add 50 µl, of deionized or sterile water.
<b>Storage</b>	Store lyophilized/reconstituted at -20 °C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please, remember to spin tubes briefly prior to opening them to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tubes.

**Application information**

<b>Recommended dilution</b>	1 : 1000 (WB)
<b>Expected   apparent MW</b>	58.4   50 kDa
<b>Confirmed reactivity</b>	<i>Arabidopsis thaliana</i>
<b>Predicted reactivity</b>	<i>Brassica oleracea</i> , <i>Brassica napus</i> , <i>Camelina sativa</i> , <i>Capsella rubella</i> , <i>Eutrema salsugineum</i> , <i>Raphanus sativus</i> Species of your interest not listed? <a href="#">Contact us</a>
<b>Not reactive in</b>	<i>Zea mays</i>
<b>Selected references</b>	To be added when available, antibody available in February 2023.



20 µg of total protein extracted freshly from 7 days-old *Arabidopsis thaliana* seedlings with Lysis buffer (50 mM Tris-HCl pH 7.5, NaCl 100 mM, 0.1% Nonidet and Protease inhibitor cocktails) and denatured with Laemmli buffer at 75°C for 5 min. The samples were separated on 10% SDS-PAGE and blotted 1h to nitrocellulose using semi-dry transfer. Blot was blocked with 1% milk for RT/ 1h with agitation. Blot was blocked in the primary antibody at a dilution of 1: 1 000 for 4°C/ON with agitation in PBS-T. The antibody solution was decanted and the blot was rinsed briefly twice, then washed once for 10 min and 3 times for 10 min in PBS-T at RT with agitation. Blot was incubated in Agrisera matching secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated [AS09 602](#)) diluted to 1: 25 000 for 1h/RT with agitation. The blot was washed as above and developed for 5 min with Agrisera [ECLBright](#). Exposure time was 10 seconds.

Courtesy of Dr. Emilio Gutierrez Beltran, University of Seville, Spain