

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no AS13 2644

SOD1 aa 58-72 | Superoxide dismutase 1, soluble (clone number 15,13)

Product information

Immunogen KLH-conjugated synthetic peptide derived from human SOD1 sequence, amino acids 58-72 P00441

Peptide used to elicit this antibody is not conserved in SOD2, 3 and 4.

Host Mouse

Clonality Monoclonal

Subclass/isotype | IgG1

Purity Immunogen affinity purified serum in PBS pH 7.4.

Format Lyophilized

Quantity 50 μg

Reconstitution For reconstitution add 50 μl of sterile water

Storage Store (vonhilized/reconstituted at -20°C; once reconstituted make aliquots to ave

Store lyophilized/reconstituted at -20 ° C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to

the cap or sides of the tube.

Application information

Recommended dilution 1:1000-1:10 000 (ELISA), 1:1000 (WB)

Expected | apparent 15.9 kDa


MW

Confirmed reactivity | Human

Predicted reactivity Bovine, Chimpanzee, Dog, Goat, Guinea Pig, Mouse, Pig, Rabbit, Rat, Schizosaccharomyces pombe, Sheep

Not reactive in No confirmed exceptions from predicted reactivity are currently known

application example



100,10 and 1 ng of recombinant human SOD1 were separated by 4-20 % SDS-PAGE and transferred electrophoretically (25V, 10 min) onto PVDF membrane. Non-specific binding sites were blocked by incubating membrane with 5 % dry milk in PBS, 0.1 % Tween 20 for 1h at room temperature (RT) with agitation. The membrane was thereafter incubated with the primary antibody SOD1 as 58-72 at a dilution of 1:1 000 for 3h at RT with agitation. The antibody solution was decanted and the membrane was inseed 3 times for 5 min in PBS-T (0.05 %) at RT with agitation. The membrane was the secondary antibody (Rabbit Anti-Mouse IgG – HRP conjugated (DAKO) at a 1:1 000 dilution) for 1h at RT with agitation. The membrane was washed as above and developed for 5 min with Amersham ECL prime western blotting detection reagent according to the manufacturer's instructions (GE Healthcare). Exposure time was 15 s.