


Product no **AS22 4820****Anti-SQE1-3 | Squalene epoxidase1-3****Product information**

<b>Immunogen</b>	KLH-conjugated peptide derived from <i>Arabidopsis thaliana</i> SQE protein sequences, UniProt: <a href="#">Q9SM02</a> , <a href="#">Q81000</a> , <a href="#">Q8VYH2</a> , , TAIR: At1g58440, At2g22830, At4g37760
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Affinity purified serum, in PBS pH 7.4
<b>Format</b>	Lyophilized
<b>Quantity</b>	50 µg
<b>Reconstitution</b>	For reconstitution, add 50 µl of sterile or deionized water.
<b>Storage</b>	Store lyophilized/reconstituted at -20 °C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please, remember to spin tubes briefly prior to opening them to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tubes.

**Application information**

<b>Recommended dilution</b>	1 : 1000 (WB)
<b>Expected   apparent MW</b>	64.7   59.5 kDa (due to N-terminal or C-terminal processing)
<b>Confirmed reactivity</b>	<i>Arabidopsis thaliana</i>
<b>Predicted reactivity</b>	<i>Brassica napus</i> , <i>Zea mays</i> , <i>Oryza sativa</i> , <i>Brachypodium distachyon</i> , <i>Triticum sp.</i> , <i>Hordeum vulgare</i> Species of your interest not listed? <a href="#">Contact us</a>
<b>Not reactive in</b>	No confirmed exceptions from predicted reactivity are currently known
<b>Selected references</b>	To be added when available, antibody available in March 2025.

**Samples:**

- 1 - 20 µg of *Withania somnifera* whole leaf extract (leaves) – adult plant, in soil, full watering
  - 2 - 20 µg of *Withania somnifera* whole leaf extract (leaves) – adult plant, in soil, drought treatment
  - 3 - 20 µg of *Withania somnifera* whole leaf extract (leaves) – adult plant, in soil, yeast extract treatment
  - 4 – No samples
  - 5 - 20 µg of *Arabidopsis thaliana* whole leaf extract (leaves) – rosette development stage, in soil
  - 6 – 20 µg of *Arabidopsis thaliana* SAIL\_1252\_E02 (insertion in G6PD1 – At5g35790) - rosette development stage, in soil
- Mark: MW markers

20 µg/well of total protein freshly extracted from *Withania somnifera* and *Arabidopsis thaliana* have been used. Exact buffer components were: TrisHCl 100 mM, MgCl<sub>2</sub> 10 mM, NaEDTA 4 mM, 10% of glycerol and 1% of protease inhibitor solution; and denatured with exact buffer components at 95 °C/5 min. Samples were separated on 10 % SDS-PAGE and blotted for 7 minutes to PVDF (pore size of 0.2 µm), using a Trans-BlotTurbo Transfer System. The blot was blocked with 5% milk for/RT with agitation. The blot was incubated in the primary antibody at a dilution of 1: 500 in TBS-T ON/4 °C with agitation. The antibody solution was decanted, and the blot was rinsed briefly twice, then washed once for 15 min and 3 times for 5 min in TBS at RT with agitation. Blot was incubated in a matching secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated) diluted to 1: 25 000 in TBS for 1 h/RT with agitation. The blot was washed as above and developed with the following



This product is **for research use only** (not for diagnostic or therapeutic use)

contact: [support@agrisera.com](mailto:support@agrisera.com)

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | [www.agrisera.com](http://www.agrisera.com)

chemiluminescent detection reagent: Agrisera ECL Bright. Exposure time was 45 seconds.  
For lower background signal, primary antibody should be incubated 1h/RT at the dilution of 1: 1000.

Courtesy of Dr. Simone Landi, Università degli Studi di Napoli Federico II, Italy