


Product no **AS11 1764****Anti-STM | Homeobox protein SHOOT MERISTEMLESS****Product information**

Immunogen	KLH-conjugated peptide chosen from <i>Arabidopsis thaliana</i> STM protein sequence, UniProt: Q38874 , TAIR: At1g62360
Host	Rabbit
Clonality	Polyclonal
Purity	Immunogen affinity purified serum in PBS pH 7.4.
Format	Lyophilized
Quantity	50 µg
Reconstitution	For reconstitution add 50 µl of sterile water
Storage	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

Application information

Recommended dilution	1 : 1000 (WB)
Expected apparent MW	42.7 kDa
Confirmed reactivity	<i>Arabidopsis thaliana</i>
Predicted reactivity	<i>Brassica napus</i>
Not reactive in	No confirmed exceptions from predicted reactivity are currently known

application example

10 µg of total protein from *Arabidopsis thaliana* seedling tissue extracted with Extractions buffers 1 or 2 (see below) were separated on 10 % SDS-PAGE and blotted 1h to PVDF using semi-dry transfer. Blots were blocked with milk for 2h at room temperature (RT) with agitation. Blot was incubated in the primary antibody at a dilution of 1: 1000 in milk overnight at 4 °C with agitation. The antibody solution was decanted and the blot was rinsed twice with milk, then washed once for 5 mins with PBS, 10 mins with 2% milk + PBST, 5 mins with PBST at RT with agitation. Blot was incubated in secondary antibody (goat anti-rabbit IgG horse radish peroxidase conjugated, from Agrisera [AS09 602](#)) diluted to 1:75 000 in for 1h at RT with agitation. The blot was washed as above and developed for 5 min with ECL according to the manufacturer's instructions. Exposure time was variable.

Extraction buffers:

- 50mM tris-acetate pH 7.9 100mM potassium acetate 20% glycerol 1mM EDTA 1mM DTT 1x Roche Protease inhibitor cocktail
- 100mM tris-cl pH 7.5 75mM NaCl 15mM EGTA 15mM MgCl₂ 1mM DTT 400 mM B glycophosphate 1x protease inhibitor cocktail NaFl, Na pyrophosphate, Na vanadate.

Courtesy Dr. Simon Scofield, Cardiff University, UK