


Product no **AS22 4844****TIP1;1, TIP1;2 | tonoplast intrinsic protein 1-1, 1-2 (gamma)****Product information**

<b>Immunogen</b>	KLH-conjugated synthetic peptide conserved in <i>Raphanus sativus</i> TIP1;1 and TIP1;2 (protein accession number available in Suga et al. 2001). Peptide is also conserved in <i>Arabidopsis thaliana</i> TIP1-1 <a href="#">P25818</a> , <a href="#">At2g36830</a> , TIP1-2 <a href="#">Q41963</a> , <a href="#">At3g26520</a>
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Antigen affinity purified serum, in PBS pH 7.4
<b>Format</b>	Lyophilized
<b>Quantity</b>	50 µg
<b>Reconstitution</b>	For reconstitution, add 50 µl of sterile water.
<b>Storage</b>	Store lyophilized/reconstituted at -20°C; make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

**Application information**

<b>Recommended dilution</b>	1 : 1000 (WB)
<b>Expected   apparent MW</b>	25,8   23 kDa
<b>Confirmed reactivity</b>	<i>Arabidopsis thaliana</i> , <i>Oryza sativa</i>
<b>Predicted reactivity</b>	<i>Brassica napus</i> , <i>Gossypium hirsutum</i> , <i>Hordeum vulgare</i> , <i>Populus trichocarpa</i> , <i>Raphanus sativus</i> , <i>Ricinus communis</i> Species of your interest not listed? <a href="#">Contact us</a>
<b>Not reactive in</b>	No confirmed exceptions from predicted reactivity are currently known
<b>Additional information</b>	Protein or membrane sample should be treated at 70°C for 10 min before loading on the gel.  Diluted antibody solution can be used 2 to 3 times within one month if it contains 0.1 % sodium azide as preservative and is stored at -20°C to -80°C.  Triton X-100 should not be included in the protein extraction buffer, when cell organelles or membrane proteins must be separated from soluble proteins. Because, Triton X breaks membrane structure and solubilizes most membranes proteins. Furthermore, it should be noted that Triton X at high concentrations binds SDS and mask the detergent effect of SDS for SDS-PAGE. Also, micelles of Triton X behave as a large complex with molecular mass of 90 kDa at high concentrations in SDS-PAGE.
<b>Selected references</b>	<a href="#">Mao</a> and Sun (2015). Arabidopsis seed-specific vacuolar aquaporins are involved in maintaining seed longevity under the control of ABSCISIC ACID INSENSITIVE 3. J Exp Bot. 2015 May 26. pii: erv244. <a href="#">Suga</a> et al. (2001). Specificity of the accumulation of mRNAs and proteins of the plasma membrane and tonoplast aquaporins in radish organs. Planta 212:294-304.



bet-mercaptoethanol). Sample was denaturized at 96 °C for 10 min. Samples were separated on 12.5% SDS-PAGE and blotted for 45 min to nitrocellulose, using: semi-dry transfer. Blot was blocked with 5 % milk or for: 1h/RT with agitation. Blot was incubated in the primary antibody at a dilution of 1: 1.000 in TBS-T for ON/4 °C with agitation. The antibody solution was decanted, and the blot was rinsed briefly twice, then washed once for 15 min and 3 times for 5 min in TBS-T at RT with agitation. Blot was incubated in matching secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated, [As09 602](#), Agrisera) diluted to 1: 25.000 in for 1 h at RT with agitation. The blot was washed as above and developed with a following chemiluminescent detection reagent: AgriseraBright ([AS16 ECL-N-10](#)).

Courtesy of Dr. Andras Bittner, IZMB - Zellbiologie der Pflanzen, University of Bonn, Germany