

Product no **AS20 4440****Anti-BirA (mutated/TurboID)****Product information**

Immunogen	Recombinant mutated BirA protein from <i>E.coli</i> produced using the following plasmid: TurboID-His6_pET21a, (Plasmid #107177). Expression was done in a vector that allowed for the generation of an untagged protein (without HIS6tag).
Host	Rabbit
Clonality	Polyclonal
Purity	Immunogen affinity purified serum in PBS pH 7.4.
Format	Lyophilized
Quantity	50 µg
Reconstitution	For reconstitution add 50 µl, of sterile water
Storage	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

Application information**Recommended dilution** 1 : 5000 (WB)**Expected | apparent MW** 35 kDa**Confirmed reactivity** BirA (mutated/TurboID)**Predicted reactivity** mini TurboID**Not reactive in** No confirmed exceptions from predicted reactivity are currently known

Selected references


[Song](#) et al. (2025). Tracking the genome-wide occupancy of Arabidopsis LEAFY COTYLEDON1 in endosperm development. *Commun Biol*. 2025 May 21;8(1):771. doi: 10.1038/s42003-025-08075-y.

[Persyn](#) et al. (2024). A Nitrogen-Specific Interactome Analysis Sheds Light on the Role of the SnRK1 and TOR Kinases in Plant Nitrogen Signaling. *Mol Cell Proteomics*. 100842. doi: 10.1016/j.mcpro.2024.100842.

[Liu](#) et al. (2024). Activation of kappa opioid receptor suppresses post-traumatic osteoarthritis via sequestering STAT3 on the plasma membrane. *Cell Commun Signal*. 2024 Jun 18;22(1):335. doi: 10.1186/s12964-024-01709-4.

[Medica](#) et al. (2023). Proximity-dependent mapping of the HCMV US28 interactome identifies RhoGEF signaling as a requirement for efficient viral reactivation. *PLoS Pathog*. 2023 Oct 2;19(10):e1011682. doi: 10.1371/journal.ppat.1011682.


[Shi](#) et al. (2023). Protocol to identify protein-protein interaction networks in Solanum tuberosum using transient TurboID-based proximity labeling. *STAR Protoc*. 2023 Sep 20;4(4):102577. doi: 10.1016/j.xpro.2023.102577.



- 1 – 20 µg *A. thaliana* – Wt Col-0 (Negative control)
- 2 – 20 µg of *A. thaliana* expressing POI-TurboID fusion (Independent line 1)
- 3 – 20 µg of *A. thaliana* expressing POI-TurboID fusion (Independent line 2)
- 4 – 20 µg of *A. thaliana* expressing POI-TurboID fusion (Independent line 3)
- 5 – 10 ng of purified TurboID (Positive control)

20 µg/well of total protein were extracted from *Arabidopsis thaliana* leaf material in diluted HENS (25mM HEPES pH 7.7, 1mM EDTA, 2.5 % SDS) and stored at -80°C. Samples were denatured in 1x protein loading dye (0.5% Sodium dodecyl Sulfate, 0.002% Bromophenol Blue, 10% glycerol, and 50 mM Tris-HCl pH6.8) at 95°C for 5 min. Samples were separated on 4-16% gradient SDS-PAGE gel and blotted 1h to a nitrocellulose membrane (pore size of 0.45 µm), using a semi-dry transfer. Blot was blocked with 5% milk in TBS-T at 4°C/ON without agitation. Blot was incubated in the primary antibody at a dilution of 1:5000 in 5% milk in TBS-T, at 4°C, ON without agitation. The antibody solution was decanted and the blot was rinsed briefly, then washed three times for 10 min in TBS-T at RT with agitation. Blot was incubated in Agrisera matching secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated, [AS09 602](#)) diluted to 1:25 000 in TBS-T for 1h at RT with agitation. The

blot was washed as above and developed for 2 min with Agrisera ECLBright. Exposure time was 172 seconds.




Samples:

- 1 - wildtype strain DDY902
- 2 - Spc105-TurboID-3xmyc, MW: 135 kDa
- 3 - Cse4-TurboID-3xmyc, MW: 57 kDa
- 4 - Ndc10-TurboID-3xmyc, MW: 140 kDa

1 µg/well of total protein extracted freshly from *Saccharomyces cerevisiae* according to [Kushnirov, 2000](#). Exact buffer components were: (6xLD = 300mM Tris pH6.8, 12% SDS, 60% Glycerol, 4% beta-Mercaptoethanol, Bromphenol blue) and denatured with 50µl 2x LD (in ddH₂O diluted) at 95°C 5 min. Samples were separated on 10 % SDS-PAGE and blotted for 2 h, 400 mA constant to nitrocellulose (pore size of 0.45 µm), using: wet transfer in the cold. Blot was blocked with 5 % milk for: 1h/RT with agitation. Following membrane washes, the blot was incubated in the primary antibody at a dilution of 1:5 000 in 5% milk in TBST with agitation ON/4°C. The antibody solution was decanted, and the blot was rinsed briefly twice, then washed once for 15 min and 3 times for 5 min in TBS-T at RT with agitation. Blot was incubated in matching secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated) diluted to 1: 20 000 in 5% milk in TBST for 1 h/RT with agitation. The blot was washed as above and developed with a following chemiluminescent detection reagent. Exposure time was 2 minutes.

Courtesy of Dr. Sonja Haberkorn, Molekulare Genetik, Universität Duisburg-Essen, Germany



1 – 1 ng of purified TurboID 2 – 5 ng of purified TurboID 3 – 10 ng of purified TurboID 4 – 25 ng of purified TurboID 5 – 50 ng of purified TurboID 6 – 75 ng of purified TurboID Mark: PageRuler™ Plus Prestained Protein Ladder; ThermoFisher Scientific; MW of TurboID = 35 kDa

1ng, 5 ng, 10 ng, 25 ng, 50 ng, and 75 ng loaded into wells 1, 2, 3, 4, 5, and 6, respectively, of purified TurboID in 1x Phosphate Buffer Saline (PBS) were stored at -80°C. Samples were denatured in 1x protein loading dye (0.5% Sodium dodecyl Sulfate, 0.002% Bromophenol Blue, 10% glycerol, and 50 mM Tris-HCl pH6.8) at 95°C for 5 min. Samples were separated on 4-16% gradient SDS-PAGE gel and blotted 1h to a nitrocellulose membrane (pore size of 0.45 µm), using a semi-dry transfer. Blot was blocked with 5% milk in TBS-T at 4°C/ON without agitation. Blot was incubated in the primary antibody at a dilution of 1:1000 in 5% TBS-T, at 4°C, ON without agitation. The antibody solution was decanted and the blot was rinsed briefly, then washed three times for 10 min in TBS-T at RT with agitation. Blot was incubated in Agrisera matching secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated, [AS09 602](#)) diluted to 1:25 000 in TBS-T for 1h at RT with agitation. The blot was washed as above and developed for 2 min with Agrisera ECLBright. Exposure time was 92 seconds.



This product is **for research use only** (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Courtesy of Eli Gordon and Dr. Patrick Treffon, Elizabeth Vierling Lab Department of Biochemistry and Molecular Biology University of Massachusetts Amherst, USA