

Product no **AS17 4158****Anti-V-ATPase holoenzyme****Product information**

Immunogen	V-ATPase complex from <i>Avena sativa</i> purified by gel filtration Ward and Sze 1992
Host	Rabbit
Clonality	Polyclonal
Purity	Serum
Format	Lyophilized
Quantity	50 µl
Reconstitution	For reconstitution add 50 µl, of sterile water
Storage	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.

Application information

Recommended dilution	1 : 400 - 1: 1000 (WB)
Expected apparent MW	100, 44 , 42, 36 , 16, 13 kDa
Confirmed reactivity	<i>Avena sativa</i>
Predicted reactivity	higher plants
Not reactive in	No confirmed exceptions from predicted reactivity are currently known
Additional information	Antibody reacts with V-ATPase subunits VHAa (100 kDa), A, B, 44 , 42, 36 , 16, 13 kDa, and with Calnexin
Selected references	Ward et al (1992).Dissociation and Reassembly of the Vacuolar H-ATPase Complex from Oat Roots. Plant Physiol. 99(1):161-9. doi: 10.1104/pp.99.1.161.